

**ROWLEY
BIOCHEMICAL
INC.**

**DANVERS INDUSTRIAL PARK
10 ELECTRONICS AVENUE,
DANVERS MA 01923
TEL: 978-739-4883 FAX: 978-739-5640
www.rowleybio.com**

F-379 HART'S METHOD FOR ELASTIC FIBERS

FIXATION: **10% Buffered Neutral Formalin**

SECTION: Paraffin, @ 6 microns

STAINING PROCEDURE:

1. Deparaffinize and hydrate to distilled water.
2. Oxidize in **Potassium Permanganate, 0.25%, Aqueous**, (F-379-3), 5 minutes. Rinse in distilled water for several minutes and then tone in **Oxalic Acid, 5%, Aqueous**, (F-379-4) until the sections are clear.
3. Wash in running tap water for several minutes and then transfer to distilled water.
4. Stain in working Resorcin-Fuchsin overnight (12 hours).
 To prepare working solution, mix just before use:
 10 ml **Resorcin Fuchsin Stock**, (F-379-1)
 100 ml 70% Alcohol
 2.0 ml Hydrochloric Acid, concentrated
5. Wash in running tap water for ten minutes and then rinse in distilled water.
6. Counterstain in **Van Gieson's Solution**, (F-379-2), 1 minute.
7. Dehydrate in 95% Alcohol, Alcohol and clear in Xylene, two changes each.
8. Mount.

RESULTS:

Elastic Fibers.....black – dark blue
Nuclei.....dark black blue
Collagen.....red-pink
Miscellaneous Tissue Elements.....yellow
(Depending on chosen counterstain)

REFERENCES:

Mallory, F.B.: Pathological Technique: New York: Hafner Publishing Co., c. 1968, 3rd Ed., p. 169.

Luna, L.G.: AFIP Manual of Histologic Staining Technique, New York: McGraw-Hill Book Co., 3rd ed., c. 1968, p. 79.