

## **F-383 GOMORI'S ONE STEP TRICHROME METHOD**

**FIXATION:** **10% BUFFERED NEUTRAL FORMALIN** (F-113)

**SECTION:** Paraffin @ 6 microns

### **STAINING PROCEDURE:**

1. Deparaffinize and hydrate to distilled water.
2. Place in **Bouin's Solution**, (F-383-4) @ 56° C for 1 hour. Wash well in running water until yellow color disappears.
3. Stain nuclei for 10 minutes in **Gomori's Chromium Hematoxylin**, (F-383-3) or **Weigert's Iron Hematoxylin**, (F-383-2A & F-383-2B). Wash in water.
4. **Gomori's Trichrome Stain with Light Green**, (F-383-1) or **Gomori's Trichrome Stain with Aniline Blue**, (F-383-1A) for 15 to 20 minutes.
5. Place in **Acetic Acid, 0.5%**, (F-383-5) for 2 minutes. If sections are too dark, differentiate in **Acetic-Phosphotungstic Acid Solution**, (F-383-6). Rinse in Distilled water.
6. Dehydrate in 95% alcohol, absolute alcohol, and clear in **Xylene**, (C-120) two changes each. Mount.

### **RESULTS:**

Muscle Fibers.....red  
Collagen.....green  
Nuclei.....blue to black

### **REFERENCES:**

Gomori, G.,: Amer. J. Clin. Path., 20:661, 1950.  
AFIP Manual of Histological Staining Method, 3<sup>rd</sup> ed., Ed. L. Luna, New York:  
McGraw - Hill Publications, c. 1968, p. 93.

### **GOMORI TRICHROME STAIN** Adapted for Stool

REAGENT REQUIRED: A. **Gomori's Trichrome Stain** (F-383-1) or (F-383-1A)  
B. **Acetic Acid, 0.2%, Aqueous**, (SO-250)

### **METHOD:**

1. Smear of stool or sections of tissue, not more than 5mm thick, prior stained in Hematoxylin followed by water rinse.
2. Reagent A for 5-20 minutes followed by water rinse.
3. Reagent B followed by water rinse.
4. Canada Balsam via usual reagents.

**NOTE: Wheatley, 1951 (Tech. Bull. 21:92) recommends this technique for protozoans in intestinal smear.**