

**ROWLEY
BIOCHEMICAL
INC.**

**DANVERS INDUSTRIAL PARK
10 ELECTRONICS AVENUE,
DANVERS MA 01923
TEL: 978-739-4883 FAX: 978-739-5640**

H-504, OIL RED IN PROPYLENE GLYCOL METHOD

FIXATION: 10% Buffered Neutral Formalin (F-113)

SECTION: Frozen

STAINING PROCEDURE:

1. Cut frozen section and collect in distilled water.
2. **Propylene Glycol, Absolute**, (H-504-6) for 2 minutes
3. **Oil Red O Solution**, (H-504-1) for one hour. If sections are mounted on glass slides before staining, allow staining process to continue overnight.
4. Differentiate in **Propylene Glycol, 85%**, (H-504-3) for one minute. Rinse in two changes of distilled water.
5. Stain for a few seconds in **Mayer's Hematoxylin**, (H-504-2) or **Harris' Hematoxylin**, (H-504-2A). Rinse in two changes of distilled water.
6. If over-stained differentiate in **Hydrochloric Acid, 5%**, (H-504-4); wash in water; neutralize in **Ammonia Water, 0.3%**, (H-504-5).
7. Wash in two changes of distilled water. Mount with **Glycerin Jelly** (M-12).

STAINING RESULTS:

Fat.....Red
Nuclei.....Blue

REFERENCES:

AFIP Manual of Histologic Staining Methods, Ed. L. Luna, 3rd ed.,
New York, McGraw-hill Publications, c. 1968, p. 140.

