

**ROWLEY
BIOCHEMICAL
INC.**

**DANVERS INDUSTRIAL PARK
10 ELECTRONICS AVENUE,
DANVERS MA 01923
TEL: 978-739-4883 FAX: 978-739-5640
www.rowleybio.com**

E-329 MODIFICATION OF MOWREY'S 1958 COLLOIDAL IRON STAIN FOR ACID MUCOPOLYSACCHARIDES

FIXATION: 10% Buffered Neutral Formalin (F-113) or 95% Alcohol. Bichromate fixatives should be avoided.

SECTION: Paraffin at 6 microns

STAINING PROCEDURE: (use control slide)

1. Deparaffinize and hydrate to distilled water.
2. Rinse in **Acetic Acid, 12%**, (E-329-6) for three minutes.
3. Place in working Colloidal Iron Solution for one hour.
Prepare working solution by mixing, just before use:

Muller's Colloidal Iron Stock ,(E-329-2)	20ml
Distilled Water	15ml
Glacial Acetic Acid	5ml
4. Rinse in **Acetic Acid, 12%**, (E-329-6) for four changes, three minutes each.
5. Place in Hydrochloric Acid-Potassium Ferrocyanide Solution for 20 minutes. Wash in tap water for five minutes, then rinse in distilled water.
Prepare Hydrochloric Acid-Potassium Ferrocyanide Solution by mixing just before use:

Potassium Ferrocyanide, 5%, Aqueous , (E-329-4)	50ml
Hydrochloric Acid, 5% , (E-329-5)	50ml
6. Counterstain with **Van Gieson's Solution**, (E-329-3) for five to seven minutes.
7. Without washing, pass quickly to 95% Alcohol, Absolute Alcohol and clear in Xylene (C-120) two changes each.
8. Mount with Permount (M-18).

RESULTS:

Acidic Mucopolysaccharides and Acidic Epithelial Mucins.....deep blue
Background depending on counterstain used.

REFERENCES:

AFIP, Manual of Histological Staining Methods, 3rd ed., Ed. L. Luna:
New York: McGraw Hill Publications, c. 1968, p. 167.